

## Compound Evaluation

### **Respiratory**

Davis RJ, Page KJ, Dos Santos Cruz GJ, Harmer DW, Munday PW, Williams SJ, *et al.* Expression and functions of the duodenal peptide secretin and its receptor in human lung. *Am J Respir Cell Mol Biol.* 2004, 31(3):302-308. [PubMed](#)

Watson ML, Grix SP, Jordan NJ, Place GA, Dodd S, Leithead J, Poll CT, Yoshimura T, Westwick J. Interleukin 8 and monocyte chemoattractant protein 1 production by cultured human airway smooth muscle cells. *Cytokine.* 1998, 10(5):346-352. [PubMed](#)

Coleman RA, Nials AT, Rabe KF, Vardey CJ, Watson N. Isolated, electrically-stimulated airway preparations : their use in determining beta-adrenoceptor agonist activity. *Pulm Pharmacol.* 1996, 9(2):107-117. [PubMed](#)

Emery N, Place GA, Dodd S, Lhermitte, David G, Lamblin G, Perini JM, Page AM, Hall RL, Roussel P. Mucous and serous secretions of human bronchial epithelial cells in secondary culture. *Am J Respir Cell Mol Biol.* 1995, 12(2):130-141. [PubMed](#)

Sheldrick RL, Rabe KF, Fischer A, Magnussen H, Coleman RA. Further evidence that tachykinin-induced contraction of human isolated bronchus is mediated only by NK2-receptors. *Neuropeptides.* 1995, 29(5):281-292. [PubMed](#)

Nials AT, Ball DI, Butchers PR, Coleman RA, Humbles AA, Johnson M, *et al.* Formoterol on airway smooth muscle and human lung mast cells: a comparison with salbutamol and salmeterol. *Eur J Pharmacol.* 1994, 251(2-3):127-135. [PubMed](#)

Nials AT, Coleman RA, Johnson M, Magnussen H, Rabe KF, Vardey CJ. Effects of beta-adrenoceptor agonists in human bronchial smooth muscle. *Br J Pharmacol.* 1993, 110(3):1112-1116. [PubMed](#)

Ball DI, Brittain RT, Coleman RA, Denyer LH, Jack D, Johnson M, *et al.* Salmeterol, a novel, long-acting beta 2-adrenoceptor agonist: characterization of pharmacological activity *in vitro* and *in vivo*. *Br J Pharmacol.* 1991, 104(3):665-671. [PubMed](#)

Sheldrick RL, Ball DI, Coleman RA. Characterisation of the neurokinin receptors mediating contraction of isolated tracheal preparations from a variety of species. *Agents Actions Suppl.* 1990, 31:205-209. [PubMed](#)

Coleman RA, Sheldrick RL. Prostanoid-induced contraction of human bronchial smooth muscle is mediated by TP-receptors. *Br J Pharmacol.* 1989, 96(3):688-692. [PubMed](#)

Coleman RA, Nials AT. Novel and versatile superfusion system. Its use in the evaluation of some spasmogenic and spasmolytic agents using guinea-pig isolated tracheal smooth muscle. *J Pharmacol Methods.* 1989, 21(1):71-86. [PubMed](#)

Coleman RA, Humphrey PP, Kennedy I, Levy GP, Lumley P. Comparison of the actions of U-46619, a prostaglandin H<sub>2</sub>-analogue, with those of prostaglandin H<sub>2</sub> and thromboxane A<sub>2</sub> on some isolated smooth muscle preparations. *Br J Pharmacol.* 1981, 73(3):773-778. [PubMed](#)

### **Inflammation**

Zhang JG, Hepburn L, Cruz G, Borman RA, Clark KL. The role of adenosine A2A and A2B receptors in the regulation of TNF-alpha production by human monocytes. *Biochem Pharmacol.* 2005, 69(6):883-889. [PubMed](#)

Houliston RA, Keogh RJ, Sugden D, Dudhia J, Carter TD, Wheeler-Jones CP. Protease-activated receptors upregulate cyclooxygenase-2 expression in human endothelial cells. *Thromb Haemost.* 2002, 88(2):321-328. [PubMed](#)

Romanska HM, Polak JM, Coleman RA, James RS, Harmer DW, Allen JC, *et al.* iNOS gene upregulation is associated with the early proliferative response of human lung fibroblasts to cytokine stimulation. *J Pathol.* 2002, 197(3):372-379. [PubMed](#)

McLaughlin F, Hayes BP, Horgan CM, Beesley JE, Campbell CJ, Randi AM. Tumor necrosis factor (TNF)-alpha and interleukin (IL)-1beta down-regulate intercellular adhesion molecule (ICAM)-2 expression on the endothelium. *Cell Adhes Commun.* 1998, 6(5):381-400. [PubMed](#)

Watson N, Bodtke K, Coleman RA, Dent G, Morton BE, Ruhlmann E, *et al.* Role of IgE in hyperresponsiveness induced by passive sensitization of human airways. *Am J Respir Crit Care Med.* 1997, 155(3):839-844. [PubMed](#)

Talpain E, Armstrong RA, Coleman RA, Vardey CJ. Characterization of the PGE receptor subtype mediating inhibition of superoxide production in human neutrophils. *Br J Pharmacol.* 1995, 114(7):1459-1465. [PubMed](#)

Chang CC, Alhasan S, Rosenspire AJ. G protein mediated signal transduction: membrane immunoglobulin associated phosphoproteins identified in octyl-beta-glucoside lysates of normal B cells. *Immunol Lett.* 1992, 32(3):193-200. [PubMed](#)

### **Gastrointestinal**

Borman RA, Wade J, Riddy D, Clark KL, Coleman RA. Effect of alosetron and tegaserod at the 5HT2B receptor in human colon. Presented at Digestive Diseases Week, 2005.

Root J, Clark KL, Borman RA. NECA-induced fluid secretion in human colonic mucosa is not mediated by adenosine A2A or A2B receptors. Presented at Digestive Diseases Week, 2004.

Borman RA, Tilford NS, Harmer DW, Day N, Ellis ES, Sheldrick RL, *et al.* 5-HT(2B) receptors play a key role in mediating the excitatory effects of 5-HT in human colon *in vitro*. *Br J Pharmacol.* 2002, 135(5):1144-1151. [PubMed](#)

Borman RA, Burleigh DE. Human colonic mucosa possesses a mixed population of 5-HT receptors. *Eur J Pharmacol.* 1996, 309(3):271-274. [PubMed](#)

Burleigh DE, Borman RA. Short-circuit current responses to 5-hydroxytryptamine in human ileal mucosa are mediated by a 5-HT4 receptor. *Eur J Pharmacol.* 1993, 241(1):125-128. [PubMed](#)

## **Oncology**

Hewes JC, Riddy D, Morris RW, Woodrooffe AJ, Davidson BR, Fuller B. A prospective study of isolated human hepatocyte function following liver resection for colorectal liver metastases: the effects of prior exposure to chemotherapy. *J Hepatol.* 2006, 45(2):263-270. [PubMed](#)

Harris SM, Mistry P, Freathy C, Brown JL, Charlton PA. Antitumour activity of XR5944 *in vitro* and *in vivo* in combination with 5-fluorouracil and irinotecan in colon cancer cell lines. *Br J Cancer.* 2005, 92(4):722-728. [PubMed](#)

Chinni SR, Alhasan SA, Multani AS, Pathak S, Sarkar FH. Pleiotropic effects of genistein on MCF-7 breast cancer cells. *Int J Mol Med.* 2003, 12(1):29-34. [PubMed](#)

Aranha O, Zhu L, Alhasan S, Wood DP, Jr., Kuo TH, Sarkar FH. Role of mitochondria in ciprofloxacin induced apoptosis in bladder cancer cells. *J Urol.* 2002, 167(3):1288-1294. [PubMed](#)

Upadhyay S, Neburi M, Chinni SR, Alhasan S, Miller F, Sarkar FH. Differential sensitivity of normal and malignant breast epithelial cells to genistein is partly mediated by p21(WAF1). *Clin Cancer Res.* 2001, 7(6):1782-9. [PubMed](#)

Glazyrin AL, Adsay VN, Vaitkevicius VK, Sarkar FH. CD95-related apoptotic machinery is functional in pancreatic cancer cells. *Pancreas.* 2001, 22(4):357-365. [PubMed](#)

Alhasan SA, Aranha O, Sarkar FH. Genistein elicits pleiotropic molecular effects on head and neck cancer cells. *Clin Cancer Res.* 2001, 7(12):4174-4181. [PubMed](#)

Li Y, Bhuiyan M, Alhasan S, Senderowicz AM, Sarkar FH. Induction of apoptosis and inhibition of c-erbB-2 in breast cancer cells by flavopiridol. *Clin Cancer Res.* 2000, 6(1):223-229. [PubMed](#)

Alhasan SA, Ensley JF, Sarkar FH. Genistein induced molecular changes in a squamous cell carcinoma of the head and neck cell line. *Int J Oncol.* 2000, 16(2):333-338. [PubMed](#)

Rosati R, Adil MR, Ali MA, Eliason J, Orosz A, Sebestyen F, *et al.* Induction of apoptosis by a short-chain neuropeptide analog in small cell lung cancer. *Peptides.* 1998, 19:1519-1523. [PubMed](#)

Thomsen LL, Lawton FG, Knowles RG, Beesley JE, Riveros-Moreno V, Moncada S. Nitric oxide synthase activity in human gynecological cancer. *Cancer Res.* 1994, 54(5):1352-1354. [PubMed](#)

Eliason JF, Kaufmann F, Tanaka T, Tsukaguchi T. Anti-proliferative effects of the arotinoid Ro 40-8757 on human cancer cell lines *in vitro*. *Br J Cancer.* 1993, 67:1293-1298. [PubMed](#)

Tevaeearai HT, Laurent PL, Suardet L, Eliason JF, Givel JC, Odartchenko N. Interactions of interferon-alpha2a with 5'-deoxy-5-fluorouridine in colorectal cancer cells *in vitro*. *Eur J Cancer.* 1992, 28(2-3):368-372. [PubMed](#)

Eliason J, Ramuz H, Kaufmann F. Human tumor cells exhibit a high degree of selectivity for stereoisomers of verapamil and quinidine. *Int J Cancer.* 1990, 46:113-117. [PubMed](#)

Eliason JF, Apro MS, Decrey D, Brink-Petersen M. Non-linearity of colony formation by human tumour cells from biopsy samples. *Br J Cancer.* 1985, 52:311-318. [PubMed](#)

## **CNS**

Watson J, Scott C, Clarke K, Riddy D, Woodrooffe A, Brown AM, Herdon HJ. Aripiprazole is a D<sub>2</sub> receptor antagonist in human striatal membranes, as measured using [<sup>35</sup>S]GTPγS binding. Presented at Neurosciences, November 2006. [Abstract](#)

Wang D, De Vivo M, Trippodi-Murphy C, Wade J, Ali M, Kogan JH, Tombaugh GC, Deng C, Dizon M, Ong VS, Murray M, Rowe WB, Sheldrick B, Lowe DA. MEM 1003, a novel L-type Ca<sup>2+</sup> channel modulator, as a potential therapeutic for Alzheimer's disease. *Annals of Neurology*, 2006, 60(5): 625-645. Presented at the 2006 annual meeting of the American Neurological Association, Chicago.

Minoru H, Takahara Y, Manning P, Brown T, Yasui M, Albala B, Kato A. Intrinsic activity of S-8510, a benzodiazepine inverse agonist, in the cortex of Alzheimer's patients. Presented at Neurosciences, November 2005. [Abstract](#)

Watson J, Roberts C, Scott C, Kendall I, Collin L, Day NC, Harries MH, Soffin E, Davies CH, Randall AD, Heightman T, Gaster L, Wyman P, Parker C, Price GW, Middlemiss DN. SB-272183, a selective 5-HT(1A), 5-HT(1B) and 5-HT(1D) receptor antagonist in native tissue. *Br J Pharmacol*. 2001, 133(6):797-806. [PubMed](#)

Day NC, Mitchell JM, Page KJ, Brown AM, Middlemiss DN, Scott C, Watson J, Price GW. Visualisation and characterization of functional 5HT receptors in the human dorsal raphe nucleus by [<sup>35</sup>S]-GTPγS autoradiography. *Eur J Neuroscience*. 2000, 12:410.

Day NC, Page KJ, Mitchell JM, Baxter GS, Brown AM, Coleman RA. Visualisation of functional CB1 cannabinoid receptors in the human CNS by [<sup>35</sup>S]-GTPγS autoradiography. *Society for Neuroscience Abstracts*. 1999, 25:1461.

Day NC, Page KJ, Mitchell JN, Baxter GS, Brown AM. Visualisation and characterisation of functional 5HT receptors in the human hippocampus by [<sup>35</sup>S]-GTPγS autoradiography. *Br J Pharmacol*. 1999, 128:7P.

## **Cardiovascular**

Maubach KA, Davis RJ, Clark DE, Fenton G, Lockey PM, Clark KL, Oxford AW, Hagan RM, Routledge C, Coleman RA. BGC20-1531, a novel, potent and selective prostanoid EP4 receptor antagonist: a putative new treatment for migraine headache. *Br J Pharmacol*. 2009, 156:316-327. [Abstract](#)

Davis RJ, Murdoch CE, Ali M, Purbrick S, Ravid R, Baxter GS, *et al*. EP4 prostanoid receptor-mediated vasodilatation of human middle cerebral arteries. *Br J Pharmacol*. 2004, 141(4):580-585. [PubMed](#)

Coleman RA, Mallett A, Sheldrick RL. EP4-receptors and cyclic AMP in pig venous smooth muscle: evidence with agonists and the EP4-antagonist, AH22921. *Adv Prostaglandin Thromboxane Leukot Res*. 1995, 23:241-246. [PubMed](#)

Coleman RA, Grix SP, Head SA, Louttit JB, Mallett A, Sheldrick RL. A novel inhibitory prostanoid receptor in piglet saphenous vein. *Prostaglandins*. 1994, 47(2):151-68. [PubMed](#)

Kathryn Bagot, Mozam Ali, Lee Dawson, Sandy Williams, Bob Sheldrick; Effects of rosiglitazone and sumatriptan on human isolated small and large coronary arteries. [Article Link](#)

### ***Ophthalmology***

Collison DJ, Coleman RA, James RS, Carey J, Duncan G. Characterization of muscarinic receptors in human lens cells by pharmacologic and molecular techniques. *Invest Ophthalmol Vis Sci*. 2000, 41(9):2633-2641. [PubMed](#)

### ***Reproductive / Genitourinary***

Bagot K, Chess-Williams R. Alpha(1A/L)-adrenoceptors mediate contraction of the circular smooth muscle of the pig urethra. *Auton Autacoid Pharmacol*. 2006, 26(4):345-53. [PubMed](#)

Baxter GS, Clayton JK, Coleman RA, Marshall K, Sangha R, Senior J. Characterization of the prostanoid receptors mediating constriction and relaxation of human isolated uterine artery. *Br J Pharmacol*. 1995, 116(1):1692-1696. [PubMed](#)

Senior J, Sangha R, Baxter GS, Marshall K, Clayton JK. *In vitro* characterization of prostanoid FP-, DP-, IP- and TP-receptors on the non-pregnant human myometrium. *Br J Pharmacol*. 1992, 107(1):215-221. [PubMed](#)

Senior J, Marshall K, Sangha R, Baxter GS, Clayton JK. *In vitro* characterization of prostanoid EP-receptors in the non-pregnant human myometrium. *Br J Pharmacol*. 1991, 102(3):747-753. [PubMed](#)

### ***ADMET / Drug Safety***

Somers GI, Lindsay N, Lowdon BM, Jones AE, Freathy C, Ho S, Woodrooffe AJM, Bayliss MK, and Manchee GR. A comparison of the expression and metabolizing activities of phase I and II enzymes in freshly isolated human lung parenchymal cells and cryopreserved human hepatocytes. *Drug Metab. Dispos*. 2007, 35:1797-1805. [PubMed](#)

Williams S, Sheldrick B, Steckler T, Gallacher D and Towart R. Effects of JNJ-17333030, naratriptan and sumatriptan on human isolated coronary arteries. Presented at Safety Pharmacology Society Meeting, September 2007 (Poster and Oral).

Roberts RA, Michel C, Coyle B, Freathy C, Cain K, Biotier E. Regulation of apoptosis by peroxisome proliferators. *Toxicol Lett*. 2004, 149(1-3):37-41. [PubMed](#)

Coleman RA, Bowen WP, Baines IA, Woodrooffe AJ, Brown AM. Use of human tissue in ADME and safety profiling of development candidates. *Drug Discovery Today*. 2001, 6(21):1116-1126. [PubMed](#)

Bowen WP, Carey JE, Miah A, McMurray HF, Munday PW, James RS, Coleman RA, Brown AM. Measurement of cytochrome P450 gene induction in human hepatocytes using quantitative real-time reverse transcriptase-polymerase chain reaction. *Drug Metab. Dispos.* 2000, 28:781-788. [PubMed](#)

Baines IA, Hickinson MM, Draper LM, Bottoms MA, Mitchell JN, Bowen WP, Brown AM. Cytotoxicity of the thiazolidinedione compounds troglitazone, rosiglitazone and pioglitazone in isolated human hepatocytes. Presented at the BPS Meeting, December 2000.